

## Catalogue of American Amphibians and Reptiles.

Flores Villela, O., H.M. Smith, E.A. Liner, and D. Chiszar. 2008. *Sceloporus jalapae*.

***Sceloporus jalapae* Günther**  
**Xalapa Spiny Lizard**

*Sceloporus graciosus*: Cope 1885:36 (part).

*Sceloporus graciosus*: Ferrari-Pérez 1886:193 (nec Yarrow).

*Sceloporus aeneus*: Ferrari-Pérez 1886:193 (nec Wiegmann).

*Sceloporus jalapae* Günther 1890:74. Type-locality, "Jalapa" (now Xalapa), [Veracruz], Mexico". Holotype, The Natural History Museum, London (BMNH) 1946.8.10.31, male (*vide* Günther 1890), collected by Höge, via F.D. Godman, date of collection unknown (not examined by authors).

*Lysioptychus jalapae*: Larsen and Tanner 1975:18.

• **CONTENT.** No subspecies are currently recognized.

• **DEFINITION.** Maximum SVL ca. 51 mm; dorsum light brown, with a broad dorsolateral light stripe, brighter in females, broken at intervals by a series of about 10 dark transverse streaks associated with paravertebral spots slightly darker than the ground color; a narrower, less conspicuous lateral light line between limbs; gular region lightly streaked and dotted with light blue; males with a lateral abdominal, dark-edged blue semeion on each side, widely separated; head scales more or less normal except for absence of postrostrals; internasals and nasals contacting rostrals; a single row of large supraoculars; dorsal scales 50–66, mean 55.6, keeled, mucronate, very weakly denticulate; lateral scales similar to dorsals, but smaller, larger than ventrals; latter notched, smooth except weakly keeled on posterior part of abdomen in some; femoral pores 17–21 on each side, separated by 0–3 scales; a pair of enlarged postanals in males.


• **DIAGNOSIS.** This species is one of the most distinctive of the genus, characterized by the combination of absence of postrostrals and postfemoral pockets, and presence of 17–21 femoral pores on each side. Its closest relative is *S. ochoterenae*, with which it forms a distinct species group as first noted by Thomas and Dixon (1976), and confirmed by Cole (1978) and Wiens and Reeder (1997). The two species share the absence of postrostrals and postfemoral pockets, but differ most conspicuously in number of femoral pores (17 or more in *S. jalapae*, fewer than 17 in *S. ochoterenae*) and in number of dorsals (50–66 vs 38–46 from interparietal to posterior margin of thighs). There is less sexual dimorphism in dorsal pattern in *S. jalapae*, and the preanals are not keeled in females. The shoulder dark spots are less evident in *S. jalapae*, and blue streaks and dots are present on the dorsum (absent in *S. ochoterenae*). Considering distribution as a character, their ranges



**FIGURE 1.** *Sceloporus jalapae* from Santa María Ixcatlán, Oaxaca, México. Photograph courtesy of Luis Canseco Márquez.

are allopatric, *S. jalapae* on Atlantic slopes, *S. ochoterenae* on Pacific slopes, and separated by some 100 km between the Izúcar de Matamoros (*S. ochoterenae*) and Zapotitlán Salinas (*S. jalapae*) regions in Puebla. The altitude range of *S. jalapae* (1000–2134 m) does not extend as low as in *S. ochoterenae* (520–2134 m), and that of the latter usually (1920 m the maximum known except for one record at 2134 m) does not extend as high as in the former.

• **DESCRIPTIONS.** Significant descriptions are given in Boulenger (1890, 1897), Günther (1890), Köhler and Heimes (2002), and Smith (1934a, 1937, 1939).



**Map.** Dots indicate localities of occurrence of *S. jalapae*. The circle indicates the type-locality.

• **ILLUSTRATIONS.** Color photographs appear in Köhler and Heimes (2002) and Canseco Márquez and Gutiérrez Mayén (2006). Head scales are illustrated in Smith (1934b, 1939). Cole (1978) illustrated the karyotype. A range map occurs in Smith (2001). Wiens and Reeder (1997) provide black-and-white drawings of skeletal elements, head scalation, and posterior lingual dentary teeth.

• **DISTRIBUTION.** A terrestrial species occurring southward from extreme northern Puebla through the eastern half of the state and the edge of adjacent Veracruz to central Oaxaca, in semiarid habitats to

evergreen forests. The record for Lake El Chico in Olson et al. (1987) is based on *S. megalepidurus*.

• **FOSSIL RECORD.** None. The name was mentioned by Wellstead (1982) in connection with fossils of other species of *Sceloporus* in Nebraska.

• **PERTINENT LITERATURE.** Topics are as follows: **anatomy** (Burststein et al. 1974; Larsen and Tanner 1974; Olson et al. 1986, 1987), **behavior** (Bussjaeger 1971; Carpenter 1978a,b, 1986; Carpenter and Ferguson 1977; Martins 1993; Martins et al. 2004; Purdue and Carpenter 1972a,b; Weins 1998), **conservation and ecology** (Bojórquez-Tapia et al. 1995; Canseco-Márquez 1996; Pelcastre Villafuerte 1991; Pelcastre Villafuerte and Flores Villela 1991; Woolrich Piña et al. 2005), **karyology** (Cole 1978; Gilboa 1974; Guillette et al. 1980; Hall 1971, 1973; Méndez-de la Cruz et al. 1998; Mink and Sites 1996), **phylogeny and systematics** (Benabib et al. 1997; Flores-Villela et al. 2000; Harmon et al. 2003; Reeder and Wiens 1996; Sites et al. 1992; Smith 1934b; Warheit et al. 1999; Wiens 1993, 1999, 2000; Wiens and Reeder 1997; Wills 1977), **reproduction** (Ramírez-Bautista 2003; Ramírez-Bautista et al. 2005). The species appears in the following **checklists, keys and similar compendia**: Bell et al. 2003; Canseco Márquez and Gutiérrez Mayén 2006; Casas Andreu et al. 1996, 2004; Eliosa León and Yañez Gómez 1994; Flores-Villela 1993; Flores-Villela and Gérez 1988, 1994; Flores-Villela et al. 1991, 1995; Frank and Ramus 1995; Gonzales Romero et al. 1991; Guillette and Smith 1985; Heimes 1993; Hutchins et al. 2003; Lara-Góngora 1986; Liner 1994, 2007; Maslin and Walker 1973; Ramírez Bautista et al. 1993; Smith 1934a, 1936, 1991; E.N. Smith 2001; Smith and Smith 1976, 1993; Smith and Taylor 1950a,b; Smith et al. 2000; and Thomas and Dixon 1976).

• **ETYMOLOGY.** The name is the possessive form of Jalapa, as the type-locality was then known (now Xalapa).

• **REMARKS.** This species, like *S. ochoterenae*, was mis-identified as *S. aeneus*, *S. graciosus* or *S. graciosus* from 1885 (Cope) to as late as 1897 (Boulenger), even though it was recognized as distinct in 1890. Hatchlings, 18 mm SVL, were seen in mid-July at 12.8 km (8 mi) SE Esperanza, Puebla.

• **ACKNOWLEDGMENTS.** We are much indebted to A. Resetar and L. Ford for checking the identity of questioned specimens, and to the curators of the collections at CAS, CM, CU, FMNH, IPN, KU, LACM, LSUMZ, MVZ, TCWC, TNHC, UAZ, UCM, UIMNH, UMMZ, UOSM, USNM, UTA and UTEP for providing information on the holdings in their museums (acronyms follow Leviton et al. 1985). C. McCarthy kindly verified the current application of the name *S. jalapae*, and gave us the catalog number of the holotype.

Bell, E.L., H.M. Smith, and D. Chiszar. 2003. An annotated list of the species-group names applied to the lizard genus *Sceloporus*. Acta Zool. Mex. (n.s.) 90:103–174.

Benabib, M., K.M. Kjer, and J.W. Sites, Jr. 1997. Mitochondrial DNA sequence-based phylogeny and the evolution of viviparity in the *Sceloporus scalaris* group (Reptilia, Squamata). Evolution 51: 1262–1275.

Bojórquez-Tapia, L.A., I. Azuara, E. Ezcurra, and O. Flores-Villela. 1995. Identifying conservation priorities in Mexico through geographic information systems and modeling. Ecol. Appl. 5:215–231.

Boulenger, G.A. 1890. First report on additions to the lizard collection in the British Museum (Natural History). Proc. Zool. Soc. Lond. 1890:77–86, pl. VIII–XI.

–. 1897. A revision of the lizards of the genus *Sceloporus*. Proc. Zool. Soc. Lond. 1897: 474–522, 1 pl.

Burststein, N., K.R. Larsen, and H.M. Smith. 1974. A preliminary survey of dermatoglyphic variation in the lizard genus *Sceloporus*. J. Herpetol. 8:359–369.

Bussjaeger, L.J. 1971. Phylogenetic significance of the comparative ethology of the *spinus* group of *Sceloporus* (Iguanidae). Ph.D. Diss., Univ. Oklahoma, Norman.

Canseco-Márquez, L. 1996. Estudio preliminar de la herpetofauna en la Cañada de Cuicatlán y Cerro Piedra Larga, Oaxaca. Bachelor's Diss. Benemerita, Univ. Autón. Puebla, Puebla, Mexico.

– and M.G. Gutiérrez Mayén. 2006. Guía de Campo de los Anfibios y Reptiles del Valle de Zapotitlán, Puebla. Soc. Herpetol. Mex., A.C. y Esc. Biol., BUAP.

Carpenter, C.C. 1978a. Ritualistic social behaviors in lizards, p. 253–267. In N.B. Greenberg and P.D. MacLean (eds.), Behavior and neurology of lizards: an interdisciplinary colloquium. U.S. DHEW Publ. No. (ADM) 77–491.

–. 1978b. Comparative display behavior in the genus *Sceloporus* (Iguanidae). Contrib. Biol. Geol. Milwaukee Pub. Mus. (18):1–71.

–. 1986. An inventory of the display-action-patterns in lizards. Smithson. Herpetol. Info. Serv. (68):1–18.

– and G.W. Ferguson. 1977. Variation and evolution of stereotyped behavior in reptiles, p. 335–403. In C. Gans and D.W. Tinkle (eds.), Biology of the Reptilia. Volume 7. New York, Academic Press.

Casas-Andreu, G., F.R. Méndez-de la Cruz, X. Aguilar-Miguel. 2004. Anfibios y Reptiles, p. 375–390. In A.J. García-Mendoza, M.d. Jesús Ordóñez, y M. Briones-Salas (eds.), Biodiversidad de Oaxaca. Inst. Biol. UNAM, Fond. Oax Conserv Nat. – WWF, México.

–, –, and J.L. Camarillo. 1996. Anfibios y reptiles de Oaxaca. Lista, distribución y conservación. Acta Zool. Mex. (n.s.) 69:1–35.

Cole, C.J. 1978. Karyotypes and systematics of the lizards in the *variabilis*, *jalapae*, and *scalaris* species groups of the genus *Sceloporus*. Amer. Mus. Novitates (2653):1–13.

- Cope, E.D. 1885. A contribution to the herpetology of Mexico. VI. A synopsis of the Mexican species of the genus *Sceloporus* Wieg. Proc. Am. Phil. Soc. 22:393–403.
- Eliosa León, H. and G. Yañes Gómez. 1994. Estudio preliminar de la herpetofauna de Zapotitlán de las Salinas, Pue. III Reunión Nac. Herpetol. Soc. Herpetol. Mexico, Chiapas:31 (abstract).
- Ferguson, G.W. 1977. Social displays in reptiles, p. 405–554. In C. Gans and D.W. Tinkle (eds.), Biology of the Reptilia, Vol. 7, Ecology and Behaviour A. Academic Press, New York.
- Ferrari-Pérez, F. 1886. Catalogue of animals collected by the Geographical and Exploring Commission of the Republic of Mexico. Proc. U.S. Natl. Mus. 9:125–199.
- Flores-Villela, O.A. 1993. Herpetofauna Mexicana: annotated list of the species of amphibians and reptiles of Mexico, recent taxonomic changes, and new species. Carnegie Mus. Nat. Hist. Spec. Publ. (17):iv + 73 p.
- and P. Gérez. 1988. Conservación en México: síntesis sobre vertebrados terrestres, vegetación y el uso del suelo. Inst. Nac. Investigaciones Recursos Nat., Xalapa, México.
- and –. 1994. Biodiversidad y conservación en México: vertebrados, vegetación y uso del suelo. Univ. Nac. Autón. México - UBIPRO, México, D.F.
- , E. Hernández García, and A. Nieto-Montes de Oca. 1991. Catálogo de anfibios y reptiles del Museo de Zoología, Facultad de Ciencias, Universidad Nacional Autónoma de México. Ser. Catálogos Mus. Zool. "Alfonso L. Herrera" (3):xii + 222 p.
- , K.M. Kjer, M. Benabib, and J.W. Sites, Jr. 2000. Multiple data sets, congruence, and hypothesis testing for the phylogeny of basal groups of the lizard genus *Sceloporus* (Squamata, Phrynosomatidae). Syst. Biol. 49:713–739.
- , F. Mendoza Quijano, and G. Gonzalez Porter (compl.). 1995. Recopilación de claves para la determinación de anfibios y reptiles de México. Publ. esp. Mus. Zool. (10):iv + 285 p.
- Frank, N. and E. Ramus. 1995. A Complete Guide to Scientific and Common Names of Reptiles and Amphibians of the World. NG Publ., Inc., Pottsville, Pennsylvania.
- Gilboa, I. 1974. Karyotypes of amphibians and reptiles: a biogeographic review, p. 91–156. In H.G. Dowling (ed.), 1974 Yearbook of Herpetology. H.I.S.S., Amer. Mus. Nat. Hist, New York.
- González-Romero, A., A. Ramírez-Bautista, and C.A. López-González. 1991. Contribución al conocimiento de los anfibios y reptiles del área de Xalapa, Veracruz, p. 377–385. In Memorias IV Simposia Sobre Fauna Silvestre General M.V. Manuel Cabrera Valtierra. UNAM Fac. Med. Zootécnica, Div. Educación Continua, México.
- Guillette, L.J., Jr., R.E. Jones, K.T. Fitzgerald, and H.M. Smith. 1980. Evolution of viviparity in the lizard genus *Sceloporus*. Herpetologica 36:201–215.
- and H.M. Smith. 1985. Cryptic species in the Mexican lizard complex, *Sceloporus aeneus*. Bull. Maryland Herpetol. Soc. 21:1–15.
- Günther, A.C.L.G. 1890. Biologia Centrali-Americana. Reptilia and Batrachia. R.H. Porter, and Dulau and Co., London.
- Hall, W.P. 1971. Chromosome evolution in the iguanid genus *Sceloporus*. Herpetol. Rev. 3:106.
- . 1973. Comparative population cytogenetics, speciation, and evolution of the iguanid lizard genus *Sceloporus*. Ph.D. Diss., Harvard Univ., Cambridge, Massachusetts.
- Harmon, L.J., J.A. Schulte, II, A. Larson, and J.B. Losos. 2003. Tempo and mode of evolutionary radiation in iguanian lizards. Science 301:961–964.
- Heimes, P. 1993. *Sceloporus aeneus* Wiegmann, 1828, ein Stachelleguane aus den zentralen Hochland von Mexico. Herpetofauna 15(86):27–34.
- Hutchins, M., J.B. Murphy, and N. Schlager (eds.). 2003. Grzimek's Animal Life Encyclopedia. 2nd ed. Vol.7. Reptiles. Gale Group, Inc., Farmington Hills, Michigan.
- Köhler, G. and P. Heimes. 2002. Stachelleguane: Lebensweise.Pflege.Zucht. Herpeton, Offenbach, Germany.
- Lara-Góngora, G. 1986. New distributional records for some Mexican reptiles and amphibians. Bull. Maryland Herpetol. Soc. 22:62–67.
- Larsen, K.R. and W.W. Tanner. 1974. Numeric analysis of the lizard genus *Sceloporus* with special reference to cranial osteology. Great Basin Nat. 34:1–41.
- and –. 1975. Evolution of sceloporine lizards (Iguanidae). Great Basin Nat. 35:1–20.
- Leviton, A.E., R.H. Gibbs, Jr., E. Heal, and C.E. Dawson. 1985. Standards in herpetology and ichthyology: Part I. Standard symbolic codes for institutional resource collections in herpetology and ichthyology. Copeia 1985:802–821.
- Liner, E.A. 1994. Scientific and common names for the amphibians and reptiles of Mexico in English and Spanish. Nombres científicos y comunes en Inglés y Español de los anfibios y reptiles de México. SSAR Herpetol. Circ. (23):v + 113 p.
- . 2007. A checklist of the amphibians and reptiles of Mexico. Occ. Pap. Mus. Nat. Sci. Louisiana St. Univ. (80):1–60.
- Martins, E.P. 1993. A comparative study of the evolution of *Sceloporus* push-up displays. Am. Nat. 142:994–1018.
- , A. Labra, M. Halloy, and J.T. Thompson. 2004. Large-scale patterns of signal evolution: an interspecific study of *Liolaemus* lizard headbob displays. Anim. Behav. 68:453–463.
- Maslin, T.P. and J.M. Walker. 1973. Variation, distribution and behavior of the lizard, *Cnemidophorus parvisocius* Zweifel (Lacertilia: Teiidae). Herpetologica 29:138–143.
- Méndez de la Cruz, F.R., M. Villagrán-Santa Cruz, and R.M. Andrews. 1998. Evolution of viviparity in the lizard genus *Sceloporus*. Herpetologica 54:521–532.
- Mink, D.G. and J.W. Sites, Jr. 1996. Species limits,



- phylogenetic relationships, and origins of viviparity in the *scalaris* complex of the lizard genus *Sceloporus* (Phrynosomatidae: Sauria). *Herpetologica* 52:551–571.
- Olson, R.E., B. Marx, and R. Rome. 1986. Descriptive dentition morphology of lizards of Middle and North America, I: Scincidae, Teiidae, and Helodermatidae. *Bull. Maryland Herpetol. Soc.* 22:97–124.
- , –, and –. 1987. Descriptive dentition of lizards of Middle and North America II: Iguanidae. *Bull. Maryland Herpetol. Soc.* 23:12–34.
- Pelcastre Villafuerte, L. 1991. Anfíbios y reptiles de Veracruz: uso del sistema de información climático-cartográfico INIREB-IBM. Tesis Biol., UNAM, México, D.F.
- and O.A. Flores-Villela. 1991. Lista de especies y localidades de recolecta de la herpetofauna de Veracruz, México. *Publ. Esp. Mus. Zool., Fac. Cienc., UNAM* (4):25–96.
- Purdue, J.R. and C.C. Carpenter. 1972a. A comparative study of the body movements of displaying males of the lizard genus *Sceloporus* (Iguanidae). *Behaviour* 41:68–81.
- and –. 1972b. A comparative study of the display motion in the iguanid genera *Sceloporus*, *Uta* and *Urosaurus*. *Herpetologica* 28:137–141.
- Ramirez-Bautista, A. 2003. Some reproductive characteristics of a tropical arid lizard assemblage from Zapotitlan Salinas, Puebla, Mexico. *Herpetol. Rev.* 34:328–331.
- , A. González-Romero, and C.A. López-González. 1993. Estudio preliminar de la herpetofauna del Municipio de Xalapa, p. 165–185. *In* I.R. López-Moreno (ed.), *Ecología Urbana Aplicada a la Ciudad de Xalapa*. Inst. Ecol., Ayuntamiento de Xalapa, Veracruz, México.
- , A.L. Ortíz-Cruz, M. Del Coro Arizmendi, and J. Campos. 2005. Reproductive characteristics of two syntopic lizard species, *Sceloporus gadoviae* and *Sceloporus jalapae* (Squamata: Phrynosomatidae), from Tehuacán Valley, Puebla, México. *West. North Amer. Nat.* 65:202–209.
- Reeder, T.W. and J.J. Wiens. 1996. Evolution of the lizard family Phrynosomatidae as inferred from diverse types of data. *Herpetol. Monogr.* (10):43–84.
- Sites, J.W., Jr., J.W. Archie, C.J. Cole, and O. Flores Villela. 1992. A review of phylogenetic hypotheses for lizards of the genus *Sceloporus* (Phrynosomatidae): implications for ecological and evolutionary studies. *Bull. Amer. Mus. Nat. Hist.* (213): 1–110.
- Smith, E.N. 2001. Species boundaries and evolutionary patterns of speciation among the malachite lizards (*formosus* group) of the genus *Sceloporus* (Squamata: Phrynosomatidae). Ph.D. Diss., Univ. Texas-Arlington.
- Smith, H.M. 1934a. Descriptions of new lizards of the genus *Sceloporus* from Mexico and southern United States. *Trans. Kansas Acad. Sci.* 37:263–285.
- . 1934b. On the taxonomic status of three species of lizards of the genus *Sceloporus* from Mexico and southern United States. *Proc. Biol. Soc. Washington* 47:121–134.
- . 1936. Descriptions of new species of lizards of the genus *Sceloporus* from Mexico. *Proc. Biol. Soc. Washington* 49:87–96.
- . 1937. A synopsis of the *scalaris* group of the lizard genus *Sceloporus*. *Occ. Pap. Mus. Zool. Univ. Michigan* (361):1–8.
- . 1939. The Mexican and Central American lizards of the genus *Sceloporus*. *Zool. Ser. Field Mus. Nat. Hist.* (26):1–397.
- . 1991. The status of the southern Mexican lizard *Sceloporus carinatus*. *Bull. Maryland Herpetol. Soc.* 27:195–200.
- , C. McCarthy, and D. Chiszar. 2000. Some enigmatic identifications in Boulenger's 1897 *Sceloporus* monograph (Reptilia: Sauria). *Bull. Maryland Herpetol. Soc.* 36:124–132.
- and R.B. Smith. 1976. Synopsis of the Herpetofauna of Mexico. Source Analysis and Index for Mexican Reptiles. Vol. III. John Johnson, North Bennington, Vermont.
- and –. 1993. Synopsis of the Herpetofauna of Mexico. Vol. VII. Bibliographic Addendum IV and Index, Bibliographic Addenda II-IV. Univ. Press of Colorado, Niwot, Colorado.
- and E.H. Taylor. 1950a. Type localities of Mexican reptiles and amphibians. *Univ. Kansas Sci. Bull.* 33:313–380.
- and –. 1950b. An annotated checklist and key to the reptiles of Mexico exclusive of the snakes. *Bull. U. S. Natl. Mus.* (199):v + 253 p.
- Thomas, R.A. and J.R. Dixon. 1976. A re-evaluation of the *Sceloporus scalaris* group (Sauria: Iguanidae). *Southwest. Nat.* 20:523–536.
- Warheit, K.I., J.D. Forman, J.B. Losos, and D.B. Miles. 1999. Morphological diversification and adaptive radiation: a comparison of two diverse lizard clades. *Evolution* 53:1226–1234.
- Wellstead, C.F. 1982. Lizards from the Lower Valentine Formation (Miocene) of northern Nebraska. *J. Herpetol.* 16:364–375.
- Wiens, J.J. 1993. Phylogenetic relationships of phrynosomatid lizards and monophyly of the *Sceloporus* group. *Copeia* 1993:287–299.
- . 1998. Testing phylogenetic methods with tree congruence: phylogenetic analysis of polymorphic morphological characters in phrynosomatid lizards. *Syst. Biol.* 47:427–444.
- . 1999. Phylogenetic evidence for multiple losses of a sexually selected character in phrynosomatid lizards. *Proc. R. Soc. Lond. B* 266:1529–1535.
- . 2000. Decoupled evolution of display morphology and display behaviour in phrynosomatid lizards. *Biol. J. Linn. Soc.* 70:597–612.
- and T.W. Reeder. 1997. Phylogeny of the spiny lizards (*Sceloporus*) based on molecular and morphological evidence. *Herpetol. Monogr.* (11):1–101.
- Wills, F.H. 1977. Distribution, geographic variation and natural history of *Sceloporus parvus* Smith (Sauria: Iguanidae). M.S. Thesis, Texas A&M University, College Station.

Woolrich Piña, G.A., L.O. López, and J.A. Lemos Espinal. 2005. *Anfibios y Reptiles del Valle de Zapotitlán*, Salinas, Puebla. Univ. Nac. Autón. Méx., Tlalnepantla y CONABIO, México, D.F.

---

**Oscar Flores-Villela**, Museo de Zoología, Facultad de Ciencias, UNAM, A.P. 70-399, México, D.F., 04510 México (ofv@hp.fciencias.unam.mx); **Hobart M. Smith**, Department of EEO Biology, University of Colorado, Boulder, CO 80309-0334 (hsmith@colorado.edu); **Ernest A. Liner**, 310 Malibou Boulevard, Houma, LA 70364-2598 (eliner@mobiletel.com); **David Chiszar**, Department of Psychology, University of Colorado, Boulder CO 80309-0345 (chiszar@clipr.colorado.edu).

Primary editor for this account, Andrew H. Price.

Published 30 April 2008 and Copyright © 2008 by the Society for the Study of Amphibians and Reptiles.

---